


Caltech Scientists Film Photons with Electrons

4D electron microscopy makes it possible to image photons of nanoscale structures and visualize their architecture



PASADENA, Calif.—Techniques recently invented by researchers at the California Institute of Technology (Caltech)—which allow the real-time, real-space visualization of fleeting changes in the structure of nanoscale matter—have been used to image the evanescent electrical fields produced by the interaction of electrons and photons, and to track changes in atomic-scale structures.

Papers describing the novel technologies appear in the December 17 issue of *Nature* and the October 30 issue of *Science*.

Four-dimensional (4D) microscopy—the methodology upon which the new techniques were based—was developed at Caltech's Physical Biology Center for Ultrafast Science and Technology. The center is directed by Ahmed Zewail, the Linus Pauling Professor of Chemistry and professor of physics at Caltech, and winner of the 1999 Nobel Prize in Chemistry.

Zewail was awarded the Nobel Prize for pioneering the science of femtochemistry, the use of ultrashort laser flashes to observe fundamental chemical reactions occurring at the timescale of the femtosecond (one-millionth of a billionth of a second). The work "captured atoms and molecules in motion," Zewail says, but while snapshots of such molecules provide the "time dimension" of chemical reactions, they don't give the dimensions of space of those reactions—that is, their structure or architecture.

Caltech Scientists Film Photons with Electrons

Published on Electronic Component News (<http://www.ecnmag.com>)

Zewail and his colleagues were able to visualize the missing architecture through 4D microscopy, which employs single electrons to introduce the dimension of time into traditional high-resolution electron microscopy, thus providing a way to see the changing structure of complex systems at the atomic scale. (See http://media.caltech.edu/press_releases/13207 [1].)

In the research detailed in the *Science* paper, Zewail and postdoctoral scholar Aycan Yurtsever were able to focus an electron beam onto a specific nanoscale-sized site in a specimen, making it possible to observe structures within that localized area at the atomic level.

In electron diffraction, an object is illuminated with a beam of electrons. The electrons bounce off the atoms in the object, then scatter and strike a detector. The patterns produced on the detector provide information about the arrangement of the atoms in the material. However, if the atoms are in motion, the patterns will be blurred, obscuring details about small-scale variations in the material.

The new technique devised by Zewail and Yurtsever addresses the blurring problem by using electron pulses instead of a steady electron beam. The sample under study—in the case of the *Science* paper, a wafer of crystalline silicon—is first heated by being struck with a short pulse of laser light. The sample is then hit with a femtosecond pulse of electrons, which bounce off the atoms, producing a diffraction pattern on a detector.

Since the electron pulses are so incredibly brief, the heated atoms don't have time to move much; this shorter "exposure time" produces a sharp image. By adjusting the delay between when the sample is heated and when the image is taken, the scientists can build up a library of still images that can be strung together into a movie.

"Essentially all of the specimens we deal with are heterogeneous," Zewail explains, with varying compositions over very small areas. "This technique provides the means for examining local sites in materials and biological structures, with a spatial resolution of a nanometer or less, and time resolution of femtoseconds."

The new diffraction method allows the structures of materials to be mapped out at an atomic scale. With the second technique—introduced in the *Nature* paper, which was coauthored by postdoctoral scholars Brett Barwick and David Flannigan—the light produced by such nanostructures can be imaged and mapped.

The concept behind this technique involves the interaction between electrons and photons. Photons generate an evanescent field in nanostructures, and electrons can gain energy from such fields, which makes them visible in the 4D microscope.

[Click Here](#) [2] for the rest of the article.

Caltech Scientists Film Photons with Electrons

Published on Electronic Component News (<http://www.ecnmag.com>)

Source URL (retrieved on 03/02/2015 - 11:57pm):

http://www.ecnmag.com/news/2009/12/caltech-scientists-film-photons-electrons?qt-video_of_the_day=0

Links:

[1] http://media.caltech.edu/press_releases/13207

[2] http://media.caltech.edu/press_releases/13310