

Atomic-scale visualization of electron pairing in iron superconductors

Brookhaven National Laboratory

Findings support magnetic pairing theory that could lead to new improved superconductors

May 3, 2012

UPTON, NY — By measuring how strongly electrons are bound together to form Cooper pairs in an iron-based superconductor, scientists at the U.S. Department of Energy's (DOE) Brookhaven National Laboratory, Cornell University, St. Andrews University, and collaborators provide direct evidence supporting theories in which magnetism holds the key to this material's ability to carry current with no resistance. Because the measurements take into account the electronic bands and directions in which the electrons are traveling, which was central to testing the theoretical predictions, this research strengthens confidence that this type of theory may one day be used to identify or design new materials with improved properties — namely, superconductors operating at temperatures far higher than today's. The findings are published in the May 4, 2012 issue of *Science*.

“In the best possible world you would be able to take this theory and plug in different chemical elements until you find a combination that should work as a superconductor at higher temperatures,” said team leader Séamus Davis, Director of the [Center for Emergent Superconductivity](#) [1] at Brookhaven and the J.G. White Distinguished Professor of Physical Sciences at Cornell University. Such materials could be used for real world, energy-saving technologies, such as zero-loss power transmission lines, without the need for expensive coolants.

Scientists have been trying to understand the mechanism underlying so-called “high-temperature” superconductivity ever since discovering materials that could carry current with no resistance at temperatures somewhat above the operating realm of conventional superconductors, which must be chilled to near absolute zero (0 kelvin, or -273° Celsius). Though still mighty chilly, these high- T_c materials' operating temperatures — some as high as 145K (-130° C) — offer hope that such materials could one day be designed to operate at room temperature.

One key to superconductivity is the formation of electron pairs. Scientists hypothesized that if these negatively charged particles have their magnetic moments pointing in opposite directions, they could overcome their mutual repulsion to join forces in so-called Cooper pairs — thus carrying current with no loss.

“Many people suspected you could take materials that naturally have alternating

Atomic-scale visualization of electron pairing in iron superconductors

Published on Electronic Component News (<http://www.ecnmag.com>)

magnetic moments on adjacent electrons — antiferromagnetic materials — and convert them into superconductors,” Davis said. But to prove this conjecture hasn’t been possible with copper-based, or cuprate, superconductors — the first high- T_c superconductors discovered starting some 25 years ago. “You can make a robust antiferromagnetic cuprate insulator, but in that state it’s hard to get the magnetic electrons to pair and then move around and make a superconductor,” Davis said.

Then, in 2008, when iron-based superconductors were discovered, the idea that magnetism plays a role in high- T_c superconductivity was revived. But determining that role was a very complex problem.

“In each iron atom there are five magnetic electrons, not just one,” Davis said. “And each, as it moves around the crystal, does so in a separate electronic band. In order to find out if the magnetic interactions between electrons are generating the superconductivity, you have to measure what’s called the anisotropic energy gap — how strongly bound together the electrons are in a pair — depending on the electrons’ directions on the different electronic bands.”

Theorists Dung-Hai Lee of the University of California at Berkeley, Peter Hirschfeld of the University of Florida, and Andrey Chubukov of the University of Wisconsin among others had developed different versions of a theory that predicts what those measurements should be if magnetism were the mechanism for superconductivity.

“It was our job to test those predictions,” Davis said. But at first, the techniques didn’t exist to make the measurements. “We had to invent them,” Davis said.

Two scientists working with Davis, Milan P. Allan of Brookhaven, Cornell, and the University of Saint Andrews (where Davis also teaches) and Andreas W. Rost of Cornell and St. Andrews — the lead authors on the paper — figured out how to do the experiments and identified an iron-based material (lithium iron arsenide) in which to test the predictions.

Their method, multi-band Bogoliubov quasiparticle scattering interference, found the “signature” predicted by the theorists:

“The strength of the ‘glue’ holding the pairs together is different on the different bands, and on each band it depends on the direction that the electrons are traveling — with the pairing usually being stronger in a given direction than at 45° to that direction,” Davis said.

“This is the first experimental evidence *direct* from the electronic structure in support of the theories that the mechanism for superconductivity in iron-based superconductors is due primarily to magnetic interactions,” he said.

The next step is to use the same technique to determine whether the theory holds true for other iron superconductors. “We and others are working on that now,” Davis said.

If those experiments show that the theory is indeed correct, the model could then

Atomic-scale visualization of electron pairing in iron superconductors

Published on Electronic Component News (<http://www.ecnmag.com>)

be used to predict the properties of other elements and combinations — and ideally point the way toward engineering new materials and higher-temperature superconductors.

This research was supported as part of the Center for Emergent Superconductivity, an Energy Frontier Research Center funded by the U.S. Department of Energy, Office of Science; the U.K. Engineering and Physical Sciences Research Council; the U.S. National Science Foundation; the Japan Society for the Promotion of Science; the Academia Sinica Research Program on Nanoscience & Nanotechnology; and a Royal Society Wolfson Research Merit Award.

DOE's Office of Science is the single largest supporter of basic research in the physical sciences in the United States, and is working to address some of the most pressing challenges of our time. For more information, please visit science.energy.gov [2].

Related Links

- [Scientific paper: "Anisotropic Energy Gaps of Iron-Based Superconductivity from Intraband Quasiparticle Interference in LiFeAs"](#) [3]

Number: 12-1410 | [BNL Media & Communications Office](#) [4]

Source URL (retrieved on 01/31/2015 - 11:22pm):

http://www.ecnmag.com/news/2012/05/atomic-scale-visualization-electron-pairing-iron-superconductors?qt-recent_content=0

Links:

[1] <http://www.bnl.gov/energy/ces/>

[2] <http://science.energy.gov/>

[3] <http://www.bnl.gov/bnlweb/pubaf/pr/>

[4] <http://www.bnl.gov/bnlweb/pubaf/medcom.asp>