

## Hubble unmask's ghost galaxies



*Astronomers used the NASA/ESA Hubble Space Telescope to unmask the dim, star-starved dwarf galaxy Leo IV. This Hubble image demonstrates why astronomers had a tough time spotting this small-fry galaxy: It is practically invisible. The image shows how the handful of stars from the sparse galaxy are virtually indistinguishable from the background. Residing 500 000 light-years from Earth, Leo IV is one of more than a dozen ultra-faint dwarf galaxies found lurking around our Milky Way galaxy. These galaxies are dominated by dark matter, an invisible substance that makes up the bulk of the Universe's mass.*

Astronomers have puzzled over why some extremely faint dwarf galaxies spotted in our Milky Way galaxy's backyard contain so few stars. The galaxies are thought to be some of the tiniest, oldest, and most pristine galaxies in the Universe. They have been discovered over the past decade by astronomers using automated computer techniques to search through the images of the Sloan Digital Sky Survey. But an international team of astronomers needed the NASA/ESA Hubble Space Telescope to help solve the mystery of why these galaxies are starved of stars, and why so few of them have been found.

Hubble views of three of these small galaxies, the Hercules, Leo IV and Ursa Major dwarf galaxies, reveal that they all started forming stars more than 13 billion years ago -- and then abruptly stopped -- all in the first billion years after the Universe

## Hubble unmask ghost galaxies

Published on Electronic Component News (<http://www.ecnmag.com>)

---

was born in the Big Bang. In fact, the extreme age of their stars is similar to Messier 92, the oldest known globular cluster [1] in the Milky Way.

"These galaxies are all ancient and they're all the same age, so you know something came down like a guillotine and turned off the star formation at the same time in these galaxies," said Tom Brown of the Space Telescope Science Institute in Baltimore, USA, the study's leader. "The most likely explanation is a process called reionisation."

The relic galaxies are evidence for a transitional phase in the early Universe that shut down star-making factories in tiny galaxies. This phase seems to coincide with the time when the first stars burned off a fog of cold hydrogen, a process called reionisation. In this period, which began in the first billion years after the Big Bang, radiation from the first stars knocked electrons off primeval hydrogen atoms, ionising the Universe's cool hydrogen gas. [2]

The same radiation that sparked universal reionisation also appears to have squelched star-making activities in dwarf galaxies, such as those in Brown's study. The small irregular galaxies were born about 100 million years before reionisation began and had just started to churn out stars at that time. Roughly 2000 light-years wide, these galaxies are the lightweight cousins of the more luminous and higher-mass star-making dwarf galaxies near our Milky Way. Unlike their higher-mass relatives, the puny galaxies were not massive enough to shield themselves from the harsh ultraviolet light. What little gas they had was stripped away as the flood of ultraviolet light rushed through them. Their gas supply depleted, the galaxies could not make new stars.

The discovery could help explain the so-called "missing satellite problem," where only a few dozen dwarf galaxies have been observed around the Milky Way while the computer simulations predict that thousands should exist. One possible explanation for the low number discovered to date is that there has been very little, or even no star formation in the smallest of these dwarf galaxies, leaving them virtually invisible.

The Sloan survey recently uncovered more than a dozen of these galaxies in our cosmic neighbourhood. These have very few stars -- only a few hundred or thousand -- but a great deal of dark matter, the underlying scaffolding upon which galaxies are built. Normal dwarf galaxies near the Milky Way contain 10 times more dark matter than the ordinary matter that makes up gas and stars, while in these so-called ultra-faint dwarf galaxies, dark matter outweighs ordinary matter by at least a factor of 100. Astronomers think the rest of the sky should contain dozens more of these ultra-faint dwarf galaxies with few stars, and the evidence for squelched star formation in the smallest of these dwarfs suggests that there may be still thousands more with essentially no stars at all.

"By measuring the star formation histories of the observed dwarfs, Hubble has supported the theoretical explanation for the paucity of such objects, according to which star formation in the smaller clumps would be shut down by reionisation," said Jason Tumlinson of the Space Telescope Science Institute, a member of the

## Hubble unmask ghost galaxies

Published on Electronic Component News (<http://www.ecnmag.com>)

---

research team.

###

The results are published in the 1 July issue of *The Astrophysical Journal Letters*.

### Notes

The Hubble Space Telescope is a project of international cooperation between ESA and NASA.

The international team of astronomers in this study consists of T. M. Brown (STScI), J. Tumlinson (STScI), M. Geha (Yale), E. N. Kirby (Cal Tech), D. A. Vandenberg (University of Victoria), R. R. Munoz (University of Chile), J. S. Kalirai (STScI), J. D. Simon (Carnegie Institution), R. J. Avila (STScI), P. Guhathakurta (UCO/Lick), A. Renzini (Osservatorio Astronomico di Padova) and, H. C. Ferguson (STScI)

[1] Globular clusters are tightly-bound spherical collections of up to a few hundred thousand stars. They are known to contain some of the most ancient stars in the Universe. In addition, globular clusters are known to have formed in single events, so all the stars in them have the same age.

[2] The period of reionisation is also the limit for how far telescopes can see: the process is what rendered the Universe's hydrogen gas transparent to ultraviolet light.

Image credit: NASA, ESA, and T. Brown (STScI)

### Links

Science paper:

[http://www.spacetelescope.org/static/archives/releases/science\\_papers/heic1211.pdf](http://www.spacetelescope.org/static/archives/releases/science_papers/heic1211.pdf) [1]

Images of Hubble:

<http://www.spacetelescope.org/images/archive/category/spacecraft/> [2]

Original release:

[http://www.eurekalert.org/pub\\_releases/2012-07/eic-hug071012.php](http://www.eurekalert.org/pub_releases/2012-07/eic-hug071012.php) [3]

**Source URL (retrieved on 12/19/2014 - 11:52am):**

[http://www.ecnmag.com/news/2012/07/hubble-unmasks-ghost-galaxies?qt-video\\_of\\_the\\_day=0](http://www.ecnmag.com/news/2012/07/hubble-unmasks-ghost-galaxies?qt-video_of_the_day=0)

### Links:

[1] [http://www.spacetelescope.org/static/archives/releases/science\\_papers/heic1211.pdf](http://www.spacetelescope.org/static/archives/releases/science_papers/heic1211.pdf)

[2] <http://www.spacetelescope.org/images/archive/category/spacecraft/>

[3] [http://www.eurekalert.org/pub\\_releases/2012-07/eic-hug071012.php](http://www.eurekalert.org/pub_releases/2012-07/eic-hug071012.php)

## **Hubble unmasks ghost galaxies**

Published on Electronic Component News (<http://www.ecnmag.com>)

---